

PROPERTIES OF CLASSICAL LIE GROUPS

WITH EMPHASIS ON DIMENSION, COMPACTNESS,
CONNECTEDNESS AND RELATIONS AMONG THE GROUPS.

JAKOB LINDBLAD BLAAVAND

APRIL 6, 2010

DEPARTMENT OF MATHEMATICAL SCIENCES
AARHUS UNIVERSITET

In this note we will focus on the fundamentals of the classical Lie Groups. There will be no proofs or argumentation, just a bunch of facts listed together for the reader to get a quick reference for the classical Lie Groups.

This note is mainly written because of the authors own inability to remember the dimensions of the classical Lie Groups, but hopefully this note will serve as an aid for others when reading mathematical literature.

Definitions of classical Lie groups

First of all we need the definitions of the classical Lie Groups that we are working with. In the following \mathbb{F} is one of the two fields \mathbb{R} or \mathbb{C} .

- $\mathrm{GL}(n, \mathbb{F}) = \{x \in \mathrm{Mat}_{n,n}(\mathbb{F}) : \det(x) \neq 0\}$
- $\mathrm{GL}^+(n, \mathbb{F}) = \{x \in \mathrm{Mat}_{n,n}(\mathbb{F}) : \det(x) > 0\}$
- $\mathrm{SL}(n, \mathbb{F}) = \{x \in \mathrm{Mat}_{n,n}(\mathbb{F}) : \det(x) = 1\}$
- $\mathrm{O}(n, \mathbb{F}) = \{x \in \mathrm{GL}(n, \mathbb{F}) : x^T x = \mathrm{Id}\}$
- $\mathrm{SO}(n, \mathbb{F}) = \{x \in \mathrm{O}(n, \mathbb{F}) : \det(x) = 1\}$
- $\mathrm{U}(n) = \{x \in \mathrm{GL}(n, \mathbb{C}) : x^* x = \mathrm{Id}\}$
- $\mathrm{SU}(n) = \{x \in \mathrm{GL}(n, \mathbb{C}) : x^* x = \mathrm{Id}, \det(x) = 1\} = \mathrm{U}(n) \cap \det^{-1}(1)$
- $\mathrm{Sp}(n, \mathbb{F}) = \{x \in \mathrm{GL}(n, \mathbb{F}) : x^T J x = J\}$ where $J = \begin{bmatrix} 0 & -\mathrm{Id} \\ \mathrm{Id} & 0 \end{bmatrix} \in \mathrm{GL}(n, \mathbb{F})$.
- $\mathrm{Sp}(n) = \mathrm{U}(2n) \cap \mathrm{Sp}(2n, \mathbb{C})$
- $\mathrm{Spin}(n)$ is the double cover of $\mathrm{SO}(n, \mathbb{R})$. For $n > 2$ it turns out that $\mathrm{Spin}(n)$ is simply connected, and in this case it coincides with the universal covering of $\mathrm{SO}(n, \mathbb{R})$.

As you see the definitions are straight forward, only the last definition of the compact symplectic group is a bit strange. Beware that there are other definitions around, but in the end they are all equivalent to this one.

The definition of the Spin-group is a bit involved, and needs a bit of explanation, which will not be given here. The definition cannot be written as a set of matrices satisfying a specific equation.

Fundamental properties of classical Lie Groups

The main part of this script is the following table. It contains all the quick facts that you need to know about a classical Lie Group.

| <i>Group</i> | <i>Lie group</i> | <i>#connected comp</i> | <i>Compact</i> | <i>Dimension</i> | π_1 | <i>Lie algebra</i> |
|----------------------|------------------|------------------------|----------------|---------------------|-------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------|
| $GL(n, \mathbb{R})$ | \mathbb{R} | 2 | no | n^2 | \mathbb{Z} | $\mathfrak{gl}(n, \mathbb{R}) = \text{End}(\mathbb{R}^n)$ |
| $GL(n, \mathbb{C})$ | \mathbb{C} | 1 | no | n^2 | \mathbb{Z} | $\mathfrak{gl}(n, \mathbb{C}) = \text{End}(\mathbb{C}^n)$ |
| $SL(n, \mathbb{R})$ | \mathbb{R} | | no | $n^2 - 1$ | $\begin{cases} 0 & n = 1 \\ \mathbb{Z} & n = 2 \\ \mathbb{Z}_2 & n > 2 \end{cases}$ | $\mathfrak{sl}(n, \mathbb{R}) = \{x \in \mathfrak{gl}(n, \mathbb{R}) : \text{Tr}(x) = 0\}$ |
| $SL(n, \mathbb{C})$ | \mathbb{C} | 1 | | $n^2 - 1$ | 0 | $\mathfrak{sl}(n, \mathbb{C}) = \{x \in \mathfrak{gl}(n, \mathbb{C}) : \text{Tr}(x) = 0\}$ |
| $O(n, \mathbb{R})$ | \mathbb{R} | 2 | | $\frac{1}{2}n(n-1)$ | | |
| $O(n, \mathbb{C})$ | \mathbb{C} | | no, $n > 1$ | $\frac{1}{2}n(n-1)$ | | |
| $SO(n, \mathbb{R})$ | \mathbb{R} | 1 | | $\frac{1}{2}n(n-1)$ | | $\mathfrak{so}(n, \mathbb{R}) = \text{skew symmetric matrices in } \mathfrak{gl}(n, \mathbb{R})$ |
| $SO(n, \mathbb{C})$ | \mathbb{C} | | no, $n > 1$ | $\frac{1}{2}n(n-1)$ | $\begin{cases} \mathbb{Z} & n = 2 \\ \mathbb{Z}_2 & n > 2 \end{cases}$ | $\mathfrak{so}(n, \mathbb{C}) = \text{skew symmetric matrices in } \mathfrak{gl}(n, \mathbb{C})$ |
| $U(n)$ | \mathbb{R} | 1 | yes | n^2 | \mathbb{Z} | $\mathfrak{u}(n) = \text{skew hermitian matrices in } \mathfrak{gl}(n, \mathbb{C})$ |
| $SU(n)$ | \mathbb{R} | 1 | yes | $n^2 - 1$ | 0 | $\mathfrak{su}(n) = \text{traceless skew hermitian matrices in } \mathfrak{gl}(n, \mathbb{C})$ |
| $Sp(2n, \mathbb{R})$ | \mathbb{R} | 1 | no | $n(2n+1)$ | \mathbb{Z} | $\mathfrak{sp}(2n, \mathbb{R}) = \{x \in \text{End}(\mathbb{R}^{2n}) : Jx + x^T J = 0\}$ |
| $Sp(2n, \mathbb{C})$ | \mathbb{C} | 1 | no | $n(2n+1)$ | 0 | $\mathfrak{sp}(2n, \mathbb{C}) = \{x \in \text{End}(\mathbb{C}^{2n}) : Jx + x^T J = 0\}$ |
| $Sp(n)$ | \mathbb{R} | 1 | yes | $n(2n+1)$ | 0 | $\mathfrak{sp}(n) = \{x \in \text{End}(\mathbb{H}^n) : x + \bar{x}^t = 0\}$ |
| $Spin(n)$ | \mathbb{R} | 1 | yes | $\frac{1}{2}n(n-1)$ | $0, n \geq 3$ | $\mathfrak{so}(n)$ |

Identities between classical groups

There are a lot of peculiar identities between the classical Lie Group – mainly between the low dimensional, though. In the following we list the most useful. The low dimensional identities are also called the accidental isomorphisms. These isomorphisms are a consequence of low dimensional isomorphisms between root systems of different families of simple Lie algebras.

- $\mathrm{GL}(n, \mathbb{C}) \cap \mathrm{SO}(2n) = \mathrm{U}(n)$
- $\mathrm{GL}(n, \mathbb{R}) \cap \mathrm{U}(n) = \mathrm{O}(n)$
- $\mathrm{GL}(n, \mathbb{R}) \simeq \mathrm{O}(n) \times \mathbb{R}^{\frac{1}{2}n(n+1)}$
- $\mathrm{GL}(n, \mathbb{C}) \simeq \mathrm{U}(n) \times \mathbb{R}^{n \cdots n}$
- $\mathrm{SL}(n, \mathbb{R}) \simeq \mathrm{SO}(n) \times \mathbb{R}^{\frac{1}{2}n(n+1)-1}$
- $\mathrm{O}(2n+1) \simeq \mathrm{SO}(2n+1) \times \mathbb{Z}_2$
- $\mathrm{O}(2n) \simeq \mathrm{SO}(2n) \times \mathbb{Z}_2$
- $\mathrm{U}(n) \simeq \mathrm{SU}(n) \times S^1$
- $\mathrm{Spin}(1) = \mathrm{O}(1)$
- $\mathrm{Spin}(2) = \mathrm{U}(1) = \mathrm{SO}(2)$
- $\mathrm{Spin}(3) = \mathrm{Sp}(1) = \mathrm{SU}(2) = S^3$
- $\mathrm{Spin}(4) = \mathrm{Sp}(1) \times \mathrm{Sp}(1)$
- $\mathrm{Spin}(5) = \mathrm{Sp}(2)$
- $\mathrm{Spin}(6) = \mathrm{SU}(4)$
- $\mathrm{SO}(3) = \mathbb{RP}^3$
- $\mathrm{U}(n) = \mathrm{SU}(n) \times \mathrm{U}(1)$

Principal bundles

The classical groups provide us with the above isomorphisms, but they also give rise to the following principal bundles.

In the following $V_k(\mathbb{R}^n)$ is the Stiefel manifold. The points in the Stiefel manifold are k -tuples of orthonormal vectors in \mathbb{R}^n . Furthermore $G_k(\mathbb{R}^n)$ is the Grassman manifold.

